

CIEN DERIVADAS RESUELTAS Y SIMPLIFICADAS

	$f(x) = 0$	$f'(x) = 0$
1	$f(x) = 0$	$f'(x) = 0$
2	$f(x) = -7$	$f'(x) = 0$
3	$f(x) = -7x$	$f'(x) = -7$
4	$f(x) = -5x + 2$	$f'(x) = -5$
5	$f(x) = x^5 - x^3 + 3$	$f'(x) = 5x^4 - 3x^2$
6	$f(x) = 2x^7 - 3x^6 + 3x^3 - 4x^2 - 7$	$f'(x) = 14x^6 - 18x^5 + 9x^2 - 8x$
7	$f(x) = \frac{x-3}{2}$	$f(x) = \frac{x}{2} - \frac{3}{2}$ $f'(x) = \frac{1}{2}$
8	$f(x) = -\frac{x^3+x-1}{2}$	$f(x) = -\frac{x^3}{2} - \frac{x}{2} + \frac{1}{2}$ $f'(x) = -\frac{3x^2}{2} - \frac{1}{2}$
9	$f(x) = -\frac{3}{2}x^3 + \frac{2}{5}x^2 - 4$	$f'(x) = -\frac{9}{2}x^2 + \frac{4}{5}x$
10	$f(x) = \frac{3}{x^2}$	$f(x) = 3 \cdot x^{-2}$ $f'(x) = -6x^{-3} = \frac{-6}{x^3}$
11	$f(x) = -\frac{2}{x^3} + \frac{3}{x^2} - 4x$	$f(x) = -2 \cdot x^{-3} + 3 \cdot x^{-2} - 4x$ $f'(x) = +6 \cdot x^{-4} - 6 \cdot x^{-3} - 4$ $f'(x) = \frac{6}{x^4} - \frac{6}{x^3} - 4$

		$f(x) = \frac{(x+1) \cdot (x-1)}{(x+1) \cdot (x+1)}$
		$f(x) = \frac{(x-1)}{(x+1)}$
12	$f(x) = \frac{x^2 - 1}{(x+1)^2}$	$f'(x) = \frac{(x-1)' \cdot (x+1) - (x-1) \cdot (x+1)'}{(x+1)^2}$ $f'(x) = \frac{(x+1) - (x-1)}{(x+1)^2} = \frac{x+1-x+1}{(x+1)^2} = \frac{2}{(x+1)^2}$ $f'(x) = \frac{2}{(x+1)^2}$
13	$f(x) = \frac{5x^4 - 3x^3}{x^5}$	$f(x) = +5\frac{x^4}{x^5} - 3\frac{x^3}{x^5}$ $f'(x) = +5x^{4-5} - 3x^{3-5}$ $f(x) = +5x^{-1} - 3x^{-2}$ $f'(x) = -5x^{-2} + 6x^{-3}$ $f'(x) = -\frac{5}{x^2} + \frac{6}{x^3}$
14	$f(x) = \sqrt{x^3}$	$f(x) = x^{\frac{3}{2}}$ $f'(x) = \frac{3}{2}x^{\frac{3}{2}-\frac{2}{2}}$ $f'(x) = \frac{3}{2}x^{\frac{1}{2}}$ $f'(x) = \frac{3}{2}\sqrt[2]{x}$

	$f(x) = \frac{1}{x^{\frac{3}{2}}}$
	$f'(x) = x^{-\frac{3}{2}}$
	$f''(x) = -\frac{3}{2}x^{-\frac{3}{2}-\frac{2}{2}}$
15	$f(x) = \frac{1}{\sqrt{x^3}}$
	$f'(x) = -\frac{3}{2}x^{-\frac{5}{2}}$
	$f''(x) = -\frac{3}{2}\frac{1}{\sqrt[2]{x^5}}$
	$f'''(x) = -\frac{3}{2}\frac{1}{x^2\sqrt[2]{x}}$
	$f(x) = x^{\frac{3}{2}} - x^{\frac{5}{3}}$
	$f'(x) = \frac{3}{2} \cdot x^{\frac{3}{2}-\frac{2}{2}} - \frac{5}{3}x^{\frac{5}{3}-\frac{3}{3}}$
16	$f(x) = \sqrt{x^3} - \sqrt[3]{x^5}$
	$f'(x) = \frac{3}{2} \cdot x^{\frac{1}{2}} - \frac{5}{3}x^{\frac{2}{3}}$
	$f''(x) = \frac{3}{2} \cdot \sqrt{x} - \frac{5}{3}\sqrt[3]{x^2}$
	$f(x) = -3x^{\frac{1}{2}} - 2x^{\frac{2}{3}}$
	$f'(x) = -\frac{3}{2}x^{\frac{1}{2}-\frac{2}{2}} - \frac{4}{3}x^{\frac{2}{3}-\frac{3}{3}}$
17	$f(x) = -3\sqrt{x} - 2\sqrt[3]{x^2}$
	$f'(x) = -\frac{3}{2} \cdot x^{-\frac{1}{2}} - \frac{4}{3}x^{-\frac{1}{3}}$
	$f''(x) = \frac{-3}{2\sqrt{x}} - \frac{4}{3\sqrt[3]{x}}$

	$f(x) = -\frac{2}{3}x^{\frac{3}{2}} - 15^{\frac{1}{2}}x^{\frac{1}{2}} - x^{\frac{5}{3}}$
18	$f(x) = -\frac{2}{3}\sqrt{x^3} - \sqrt{15x} - \sqrt[3]{x^5}$
	$f'(x) = -\frac{2}{3} \cdot \frac{3}{2}x^{\frac{3}{2}-\frac{2}{2}} - \sqrt{15} \frac{1}{2}x^{\frac{1}{2}-\frac{2}{2}} - \frac{5}{3}x^{\frac{5}{3}-\frac{3}{3}}$
	$f'(x) = -x^{\frac{1}{2}} - \sqrt{15} \frac{5}{3}x^{-\frac{1}{2}} - x^{\frac{2}{3}}$
	$f'(x) = -\sqrt{x} - \frac{\sqrt{15}}{2\sqrt{x}} - \frac{5}{3}\sqrt[3]{x^2}$
19	$f(x) = -\frac{3}{2}\sqrt{x^3} - 2x^5 - 5x^2$
	$f'(x) = -\frac{3}{2} \cdot \frac{3}{2}x^{\frac{1}{2}-1} - 10x^4 - 10x$
	$f'(x) = -\frac{9}{4}\sqrt{x} - 10x^4 - 10x$
20	$f(x) = \frac{\sqrt{x^3}\sqrt{x}}{2\sqrt{x}} = \frac{x^{\frac{1}{2}}x^{\frac{1}{3}}}{x^{\frac{1}{2}}} = \frac{x^{\frac{1}{2}+\frac{1}{3}}}{x^{\frac{1}{2}}} = \frac{x^{\frac{5}{6}}}{x^{\frac{1}{2}}}$
	$f(x) = x^{\frac{5}{6}-\frac{1}{2}} = x^{\frac{2}{3}}$
	$f'(x) = \frac{2}{6} \cdot x^{-\frac{4}{6}} = \frac{2}{6} \cdot x^{-\frac{2}{3}}$
	$f'(x) = \frac{1}{3\sqrt[3]{x^2}}$

	$f(x) = \frac{2\sqrt{x} + \sqrt[3]{x}}{\sqrt[2]{x}} = \frac{2\sqrt{x}}{\sqrt[2]{x}} + \frac{\sqrt[3]{x}}{\sqrt[2]{x}}$	
21	$f(x) = \frac{2x^{\frac{1}{2}}}{x^{\frac{1}{2}}} + \frac{x^{\frac{1}{3}}}{x^{\frac{1}{2}}}$	$f(x) = 2 + x^{\frac{1}{6}}$
		$f'(x) = -\frac{1}{6}x^{-\frac{5}{6}}$
		$f'(x) = -\frac{1}{6x^{\frac{5}{6}}\sqrt{x}}$
22	$f(x) = (x^5 - x^3 + 3)^4$	$f'(x) = 4 \cdot (x^5 - x^3 + 3)^3 \cdot (5x^4 - 3x^2)$
		$f'(x) = 2 \cdot (x^2 - 2) \cdot (2x)$
23	$f(x) = (x^2 - 2)^2$	$f'(x) = 4x \cdot (x^2 - 2)$
		$f'(x) = 4x^3 - 8x$
24	$f(x) = (x - 1) \cdot (x + 1)^2$	$f'(x) = (x + 1)^2 + (x - 1) \cdot 2 \cdot (x + 1)$
		$f'(x) = x^2 + 2x + 1 + 2x^2 - 2$
		$f'(x) = 3x^2 + 2x - 1$
25	$f(x) = (x^5 - x^3 + 3)^4$	$f(x) = (x^5 - x^3 + 3)^4$
		$f'(x) = 4 \cdot (x^5 - x^3 + 3)^3 \cdot (5x^4 - 3x^2)$

26	$f(x) = \sqrt{(x^5 - x^3 + 3)}$ $f'(x) = \frac{1}{2} (x^5 - x^3 + 3)^{\frac{1}{2}} \cdot (5x^4 - 3x^2)$ $f'(x) = \frac{(5x^4 - 3x^2)}{2\sqrt{x^5 - x^3 + 3}}$
27	$f(x) = \sqrt[5]{x^5 - x^3 + 3}$ $f'(x) = \frac{1}{5} (x^5 - x^3 + 3)^{-\frac{4}{5}} \cdot (5x^4 - 3x^2)$ $f'(x) = \frac{(5x^4 - 3x^2)}{5\sqrt[5]{(x^5 - x^3 + 3)^4}}$
28	$f(x) = (x^5 - x^3 + 3)^{-\frac{1}{5}}$ $f'(x) = -\frac{1}{5} (x^5 - x^3 + 3)^{-\frac{6}{5}} \cdot (5x^4 - 3x^2)$ $f'(x) = -\frac{(5x^4 - 3x^2)}{5\sqrt[5]{(x^5 - x^3 + 3)^6}} = -\frac{(5x^4 - 3x^2)}{5(x^5 - x^3 + 3)^5 \sqrt[5]{(x^5 - x^3 + 3)}}$

	$f(x) = \left(\frac{x^5 - x^3 + 3}{x^2} \right)^{\frac{1}{3}}$
29	$f'(x)$ $= \frac{1}{3 \sqrt[3]{\left(\frac{x^5 - x^3 + 3}{x^2} \right)^2}} \cdot \frac{(5x^4 - 3x^2)x^2 - (x^5 - x^3 + 3) \cdot 2x}{x^4}$ $= \frac{1}{3 \sqrt[3]{\left(\frac{x^5 - x^3 + 3}{x^2} \right)^2}} \cdot \frac{(3x^5 - x^3 - 6)}{x^3}$
30	$f(x) = \sqrt[5]{\frac{x^2+x}{x+1}} = \sqrt[5]{\frac{x \cdot (x+1)}{x+1}} = \sqrt[5]{x}$ $f'(x) = \frac{1}{5 \sqrt[5]{x^4}}$
31	$f(x) = \sqrt{\frac{x^2+2x+1}{x^2-1}} = \sqrt{\frac{(x+1)(x+1)}{(x+1)(x-1)}} = \sqrt{\frac{(x+1)}{(x-1)}}$ $f'(x) = \frac{1}{2 \sqrt{\frac{(x+1)}{(x-1)}}} \frac{x-1-x-1}{(x-1)^2} = \frac{1}{\sqrt{\frac{(x+1)}{(x-1)}}} \frac{-2}{(x-1)^2} = \frac{1}{\sqrt{x^2-1}} \frac{-2}{(x-1)^2}$ $f'(x) = \frac{-1}{\sqrt{x^2-1} \cdot (x-1)}$

	$f(x) = \sqrt{\frac{x^2-1}{x^2-2x+1}} = \sqrt{\frac{(x+1).(x-1)}{(x-1).(x-1)}} = \sqrt{\frac{(x+1)}{(x-1)}}$	
32	$f(x) = \sqrt{\frac{x^2-1}{x^2-2x+1}}$	$f'(x) = \frac{1}{2\sqrt{(x+1)}} \frac{x-1-x-1}{(x-1)^2} = -\frac{1}{\sqrt{(x+1)}} \frac{1}{(x-1)^2} = -\frac{1}{\sqrt{x^2-1}} \frac{1}{(x-1)}$
		$f'(x) = \frac{-1}{\sqrt{x^2-1} \cdot (x-1)}$
33	$f(x) = e^{x+1}$	$f'(x) = e^{x+1}$
34	$f(x) = -3 \cdot e^{x+1}$	$f'(x) = -3 \cdot e^{x+1}$
35	$f(x) = 7 \cdot e^{x^2+1}$	$f'(x) = 7 \cdot e^{x^2+1} \cdot 2x = 14x \cdot e^{x^2}$
36	$f(x) = -3 \cdot e^{x^2+x-1}$	$f'(x) = -3 \cdot (2x+1) e^{x^2+x-1}$
37	$f(x) = \sqrt{e^x}$	$f'(x) = \frac{e^x}{2\sqrt{e^x}}$
38	$f(x) = \sqrt{3e^{x+1}}$	$f'(x) = \frac{3e^{x+1}}{2\sqrt{3e^{x+1}}}$
39	$f(x) = -\frac{2}{\sqrt{e^x}}$	$f(x) = -2 \cdot (e^x)^{-\frac{1}{2}}$ $f'(x) = \frac{+2}{2} \cdot (e^x)^{-\frac{3}{2}} \cdot e^x$ $f'(x) = \frac{1}{\sqrt{e^x}}$
40	$f(x) = e^{x+1} - 3e^x + 2e^{x^3}$	$f'(x) = e^{x+1} - 3e^x + 6x^2 e^{x^3}$
41	$f(x) = 3^{2x+1}$	$f'(x) = 3^{2x+1} \cdot \ln 3 \cdot 2$
42	$f(x) = 7^{x-1}$	$f'(x) = 7^{x-1} \cdot \ln 7$

43	$f(x) = 7^{x^2-1}$	$f'(x) = 7^{x^2-1} \cdot \ln 7 \cdot 2x$
44	$f(x) = -\frac{1}{\sqrt{2^x}}$	$f'(x) = -\frac{1}{2} (2^x)^{-\frac{1}{2}} \ln 2 \cdot 2^x$ $f'(x) = \frac{\ln 2}{2\sqrt{2^x}}$
45	$f(x) = 2^{x+1} - 3 \cdot 5^x$	$f'(x) = 2^{x+1} \cdot \ln 2 - 3 \cdot (5^x \cdot \ln 5)$
46	$f(x) = (2^{x+1} - 3 \cdot 5^x)^3$	$f'(x) = 3 \cdot (2^{x+1} - 3 \cdot 5^x)^2 \cdot (2^{x+1} \cdot \ln 2 - 3 \cdot (5^x \cdot \ln 5))$
47	$f(x) = \sqrt{3^{x+1}}$	$f(x) = (3^{x+1})^{\frac{1}{2}}$ $f'(x) = \frac{1}{2} \cdot (3^{x+1})^{-\frac{1}{2}} \cdot 3^{x+1} \cdot \ln 3$ $f'(x) = \frac{3^{x+1} \ln 3}{2\sqrt{3^{x+1}}}$
48	$f(x) = 7^{\sqrt{x+1}}$	$f'(x) = 7^{\sqrt{x+1}} \cdot \ln 7 \cdot \frac{1}{2\sqrt{x+1}}$
49	$f(x) = \frac{e^{3x} + e^{x^2}}{3}$	$f'(x) = \frac{e^{3x} \cdot 3}{3} + \frac{e^{x^2} \cdot 2x}{3}$ $f'(x) = e^{3x} + \frac{e^{x^2} \cdot 2x}{3}$

50	$f(x) = \frac{7^{x^2}}{x^3}$ $f'(x) = \frac{7^{x^2} \cdot \ln 7 \cdot 2x \cdot x^3 - 7^{x^2} \cdot 3x^2}{x^6}$
51	$f(x) = \frac{e^{x^2}}{x^3}$ $f'(x) = \frac{e^{x^2} \cdot 2x \cdot x^3 - e^{x^2} \cdot 3x^2}{x^6} = \frac{x^2(e^{x^2} \cdot 2 \cdot x^2 - e^{x^2} \cdot 3)}{x^6} =$ $\frac{(e^{x^2} \cdot 2 \cdot x^2 - e^{x^2} \cdot 3)}{x^4}$ $f'(x) = \frac{e^{x^2} \cdot (2 \cdot x^2 - 3)}{x^4}$
52	$f(x) = \sqrt{\frac{7^{x^2}}{x^3}}$ $f'(x) = \frac{1}{2\sqrt{\frac{7^{x^2}}{x^3}}} \cdot \frac{7^{x^2} \cdot \ln 7 \cdot 2x \cdot x^3 - 7^{x^2} \cdot 3x^2}{x^6}$
53	$f(x) = \ln(x + 3)$ $f'(x) = \frac{1}{x+3}$
54	$f(x) = 7x + \ln(x - 3)$ $f'(x) = 7 + \frac{1}{x-3}$
55	$f(x) = \ln(x^2 - 3x + 2)$ $f'(x) = \frac{1}{x^2 - 3x + 2} \cdot (2x - 3)$
56	$f(x) = \frac{1}{\ln(x-1)}$ $f'(x) = \frac{-\frac{1}{(x-1)}}{(\ln(x-1))^2} = -\frac{1}{(x-1)((\ln(x-1))^2)}$

	$f(x) = \ln \sqrt{\frac{x^2-1}{x^2-2x+1}} =$ $\ln \sqrt{\frac{(x-1).(x+1)}{(x-1).(x-1)}} = \ln \sqrt{\frac{(x+1)}{(x-1)}}$ $f'(x) = \frac{1}{\sqrt{\frac{(x+1)}{(x-1)}}} \cdot \frac{1}{2\sqrt{\frac{(x+1)}{(x-1)}}} \frac{x-1-x-1}{(x-1)^2} = ..$ $\frac{-2}{\frac{(x+1)}{(x-1)} \cdot 2 \cdot (x-1)^2} = -\frac{1}{x^2-1}$ $f'(x) = -\frac{1}{x^2-1}$
57	$f(x) = \ln \left(\sqrt{(x^5 - x^3 + 3)} \right)$ $f'(x) = \frac{1}{\sqrt{(x^5 - x^3 + 3)}} \cdot \frac{1}{2\sqrt{(x^5 - x^3 + 3)}} (5x^4 - 3x^2)$ $f'(x) = \frac{5x^4 - 3x^2}{(x^5 - x^3 + 3)}$
58	$f(x) = \ln \left(\frac{e^x - 1}{e^x + 1} \right)$ $f'(x) = \frac{1}{\frac{e^x - 1}{e^x + 1}} \cdot \frac{e^x \cdot (e^x + 1) - e^x \cdot (e^x - 1)}{(e^x + 1)^2}$ $f'(x) = \frac{+2e^x}{e^{2x} - 1}$
59	$f(x) = \log_3(x + 2)$ $f'(x) = \frac{u'}{u} \log_a e$ $f'(x) = \frac{1}{x + 2} \log_3 e$
60	$f(x) = \log(x - 3)^2$ $f'(x) = \frac{2 \cdot (x-3)}{(x-3)^2 \cdot \ln 10}$
61	$f(x) = \operatorname{sen}(x + 1)$ $f'(x) = \cos(x + 1)$
62	

63	$f(x) = \operatorname{sen}(2x^3 + 2x^2)^2$	$f'(x) = 2 \cdot \sin(2x^3 + 2x^2) \cdot \cos(2x^3 + 2x^2) (6x^2 + 4x)$
64	$f(x) = \operatorname{sen}(x + 1) + 5x$	$f'(x) = \cos(x + 1) + 5$
65	$f(x) = \sqrt{\operatorname{sen}(x + 1)}$	$f(x) = \sin(x + 1)^{\frac{1}{2}}$ $f'(x) = \frac{1}{2} \sin(x + 1)^{\frac{-1}{2}} \cdot \cos(x + 1)$ $f'(x) = \frac{\cos(x+1)}{2\sqrt{\sin(x+1)}}$
66	$f(x) = \cos(3x + 3)$	$f'(x) = -\sin(3x + 3) \cdot 3$ $f'(x) = -3 \sin(3x + 3)$
67	$f(x) = \cos(3x^2 + 3x)$	$f'(x) = -\sin(3x^2 + 3x) \cdot (6x + 3)$ $f'(x) = -3 \cdot \sin(3x^2 + 3x) \cdot (3x + 1)$
68	$f(x) = \frac{1}{\operatorname{sen}(x+1)}$	$f'(x) = -\frac{\cos(x+1)}{(\operatorname{sen}(x+1))^2}$
69	$f(x) = \frac{1}{\cos x} + \frac{1}{\operatorname{sen}(x+1)}$	$f'(x) = \frac{\operatorname{sen} x}{(\cos x)^2} - \frac{\cos(x+1)}{(\operatorname{sen}(x+1))^2}$
70	$f(x) = \frac{1}{\operatorname{sen} x} - \frac{1}{\cos(x-1)}$	$f'(x) = \frac{-\cos x}{(\operatorname{sen} x)^2} - \frac{\operatorname{sen}(x+1)}{(\cos(x-1))^2}$
71	$f(x) = \sqrt[3]{\cos(3x + 3)}$	$f'(x) = \frac{1}{3\sqrt[3]{(\cos(3x+3))^2}} \cdot -\sin(3x + 3) \cdot 3$ $f'(x) = -\frac{\operatorname{sen}(3x+3)}{3\sqrt[3]{(\cos(3x+3))^2}}$

	$f(x) = \frac{1}{\sin(x+1)} + (x^5 - x^3 + 3)^4$	$f'(x) = -\frac{\cos(x+1)}{(\sin(x+1))^2} + 4 \cdot (x^5 - x^3 + 3)^3 \cdot (5x^4 - 3x^2)$
72		
73	$f(x) = \ln(x-1) + e^{x+1}$	$f'(x) = \frac{1}{x-1} + e^{x+1}$
74	$f(x) = e^{x-3} + \cos(x+1) - x^2$	$f'(x) = e^{x-3} - \sin(x+1) - 2x$
75	$f(x) = \tan(x-5)$	$f'(x) = \sec^2(x-5)$
76	$f(x) = \tan(x^3 + 3)$	$f'(x) = \sec^2(x^3 + 3) \cdot 3x^2$
77	$f(x) = -\tan(-5x^2 - 7)$	$f'(x) = -\sec^2(-5x^2 - 7) \cdot (-10x)$ $f'(x) = \sec^2(-5x^2 - 7) \cdot 10x$
78	$f(x) = \frac{1}{\tan(x-5)}$	$f'(x) = \frac{-\sec^2(x-5)}{(\tan(x-5))^2}$
79	$f(x) = -\frac{3}{\tan(x+2)}$	$f'(x) = \frac{3 \cdot \sec^2(x+2)}{(\tan(x+2))^2}$
80	$f(x) = \sqrt{\tan(x-5)}$	$f'(x) = \frac{\sec^2(x-5)}{2 \cdot \sqrt{\tan(x-5)}}$
81	$f(x) = \arcsen(x^2 - 3)$	$f'(x) = \frac{2x}{\sqrt{1 - (x^2 - 3)^2}}$
82	$f(x) = 3x + \arcsen(3x^3 + 3x - 7)$	$f'(x) = 3 + \frac{9 \cdot x^2 + 3}{\sqrt{1 - (3x^3 + 3x - 7)^2}}$

83

$$f(x) = \arcsen \sqrt{x^2 - 3}$$

$$\begin{aligned} f'(x) &= \frac{\frac{1}{2\sqrt{x^2-3}} \cdot 2x}{\sqrt{1-(x^2-3)}} \\ &= \frac{x}{\sqrt{(-x^2+4).(x^2-3)}} \end{aligned}$$

84

$$f(x) = \arcsen \left(\frac{x+1}{x-1} \right)$$

$$\begin{aligned} f'(x) &= \frac{1}{\sqrt{1-\left(\frac{x+1}{x-1}\right)^2}} \cdot \frac{x-1-(x+1)}{(x-1)^2} \\ &= \frac{1}{\sqrt{x^2-2x+1-x^2-2x-1}} \cdot \frac{-2}{(x-1)^2} \end{aligned}$$

$$f'(x) = \frac{-2}{2 \cdot (x-1) \cdot \sqrt{-x}}$$

$$f'(x) = \frac{-1}{(x-1) \cdot \sqrt{-x}}$$

85

$$f(x) = \sqrt[3]{\sen(x^2 + 3)}$$

$$f(x) = (\sen(x^2 + 3))^{\frac{1}{3}}$$

$$f'(x) = \frac{\cos(x^2 + 3) \cdot 2x}{3 \cdot \sqrt[3]{(\sen(x^2 + 3))^2}}$$

86

$$f(x) = \sqrt[3]{\tan e^x}$$

$$f'(x) = \frac{\sec^2(e^x) \cdot e^x}{3 \cdot \sqrt[3]{(\tan(e^x))^2}}$$

87

$$f(x) = x^2 \cdot \tan \sqrt{x}$$

$$f'(x) = 2x \cdot \tan \sqrt{x} + x^2 \cdot \sec^2 \sqrt{x} \cdot \frac{1}{2\sqrt{x}}$$

88	$f(x) = \frac{1 + \sin^2 x}{x}$	$f'(x) = \frac{-2 \cdot \sin x \cdot \cos x}{x^2}$
89	$f(x) = \ln(\sin x)$	$f'(x) = \frac{1}{\sin x} \cdot \cos x$
90	$f(x) = \arctg(x^2 - 3)$	$f'(x) = \frac{1}{1 + (x^2 - 3)^2} \cdot 2x$
91	$f(x) = e^{x^2} \cdot 3 \ln(\sin x)$	$f'(x) = e^{x^2} \cdot 2x \cdot 3 \frac{1}{(\sin x)} \cdot \cos x$
92	$f(x) = e^{x+3} + \ln(x-5) \cdot \cot(x)$	$f'(x) = e^{x+3} + \frac{1}{x-5} + \operatorname{cosec}^2(x)$
93	$f(x) = \arctg(\ln x)$	$f'(x) = \frac{1}{1 + (\ln x)^2} \cdot \frac{1}{x}$
94	$f(x) = \ln(\ln x)$	$f'(x) = \frac{1}{\ln x} \cdot \frac{1}{x}$
95	$f(x) = \ln(\ln x) + \arctg(x^3 - 1)$	$f'(x) = \frac{1}{\ln x} \cdot \frac{1}{x} + \frac{1}{1 + (x^3 - 1)^2} \cdot 3x^2$
96	$f(x) = \cot(x^3 - 1)$	$f'(x) = -3x^2 \cdot \operatorname{cosec}^2(x^3 - 1)$
97	$f(x) = \sec x - e^x$	$f'(x) = \sec x \cdot \operatorname{tg} x - e^x$
98	$f(x) = \operatorname{cosec} x + \frac{x^3}{3}$	$f'(x) = -\operatorname{cosec} x \cdot \operatorname{tg} x + x^2$
99	$f(x) = \cot(x+1)$	$f'(x) = -\operatorname{cosec}^2(x+1)$
100	$f(x) = e^{x^2} \cdot \cot(x^3 - 1)$	$f'(x) = e^{x^2} \cdot 2x + 3x^2 \cdot \operatorname{cosec}^2(x^3 - 1)$